

Bleeding risk differences after TAVR according to the ARC-HBR criteria: insights from SCOPE 2

Philippe Garot^{1*}, MD; Antoinette Neylon¹, MD; Marie-Claude Morice¹, MD; Corrado Tamburino², MD, PhD; Sabine Bleiziffer³, MD; Holger Thiele⁴, MD; Smita Scholtz⁵, MD; Rene Schramm³, MD, PhD; James Cockburn⁶, MD; Michael Cunningham⁷, MD; Alexander Wolf⁸, MD; Marco Barbanti⁹, MD; Didier Tchétché¹⁰, MD; Paolo Pagnotta¹¹, MD; Martine Gilard¹², MD; Francesco Bedogni¹³, MD; Eric Van Belle¹⁴, MD; Mariuca Vasa-Nicotera¹⁵, MD; Alaide Chieffo¹⁶, MD; Kris Bogaerts¹⁷, PhD; Christian Hengstenberg¹⁸, MD; Davide Capodanno², MD, PhD

The authors' affiliations can be found in the Appendix paragraph.

GUEST EDITOR: Franz-Josef Neumann, MD; Department of Cardiology and Angiology II, University Heart Center Freiburg - Bad Krozingen, Bad Krozingen, Germany

This paper also includes supplementary data published online at: <https://eurointervention.pronline.com/doi/10.4244/EIJ-D-21-01048>

KEYWORDS

- bleeding
- TAVR

Abstract

Background: The Academic Research Consortium – High Bleeding Risk (ARC-HBR) initiative defined conditions associated with percutaneous coronary intervention (PCI)-related bleeding.

Aims: We sought to further explore these HBR conditions in the setting of transcatheter aortic valve replacement (TAVR).

Methods: Patients from the SCOPE 2 trial were stratified by their bleeding risk status based on the ARC-HBR definitions. Baseline and procedural characteristics, as well as key clinical outcomes including Bleeding Academic Research Consortium (BARC) 3-5 bleeding, were compared in ARC-HBR positive (HBR+) and ARC-HBR negative (HBR-) patients.

Results: Of 787 patients randomised in SCOPE 2 and included in this study, 633 were HBR+ (80.4%). Compared with HBR- patients, those HBR+ were older and more frequently presented with diabetes, a history of coronary artery disease, atrial fibrillation, prior cerebrovascular accident, and a Society of Thoracic Surgeons predicted risk of 30-day mortality (STS-PROM) ($4.9 \pm 2.9\%$ vs $3.3\% \pm 2.1\%$; $p < 0.0001$). In addition, HBR+ patients were more frequently on oral anticoagulation therapy. At 1 year, HBR+ patients had higher rates of all-cause death (12.4% vs 4.3%, respectively, risk difference 8.09%; 95% confidence interval [CI]: 3.76-12.41; $p = 0.0002$); the rates of BARC 3-5 type bleeding were relatively high but not statistically different compared with HBR- patients (7.7% vs 6.1%, risk difference 1.67%; 95% CI: -2.72 to 6.06; $p = 0.46$). Subgroup analyses for bleeding events showed no significant interaction in terms of STS-PROM score, age, or medications.

Conclusions: The ARC-HBR criteria failed to isolate a subgroup of patients at higher bleeding risk in TAVR patients from a randomised trial. These findings have potential implications, especially for the selection of post-TAVR antithrombotic regimens based on individual bleeding-risk profiles. Specific HBR criteria should be defined for TAVR patients.

*Corresponding author: Institut Cardiovasculaire Paris-Sud, Hôpital Jacques Cartier, 6 avenue du Noyer Lambert, 91300 Massy, France. E-mail: pgarot@angio-icps.com

Abbreviations

ARC	Academic Research Consortium
BARC	Bleeding Academic Research Consortium
DAPT	dual antiplatelet therapy
HBR	high bleeding risk
OAC	oral anticoagulant
PCI	percutaneous coronary intervention
SAVR	surgical aortic valve replacement
STS-PROM	Society of Thoracic Surgeons – predicted risk of mortality
TAVR	transcatheter aortic valve replacement

Introduction

Transcatheter aortic valve replacement (TAVR) has proven to be an effective and minimally invasive procedure in patients suffering from severe aortic stenosis^{1,2}. TAVR is an alternative to surgical aortic valve replacement (SAVR) for a large proportion of patients with severe aortic stenosis, especially for those who are older or present with intermediate or high risk for surgery^{3,4}. Since the late 2000's, TAVR has led to constant improvement in clinical outcomes with the development of techniques and technological ameliorations along with increased operator experience. Although less prevalent compared with SAVR, the risk of major bleeding after TAVR has been estimated to be as high as approximately 6% and is associated with a three-fold increase in one-year mortality⁵⁻⁷.

Unlike percutaneous coronary intervention (PCI)-related bleeding risk conditions that have been recently defined by an Academic Research Consortium - High Bleeding Risk (ARC-HBR) consensus document⁸, those associated with major bleeding remain insufficiently explored after TAVR. The Valve Academic Research Consortium (VARC)-3 consensus provided an overview of risk assessment after TAVR that included the definitions of bleeding, but the conditions that increased this risk were not the subject of this initiative⁹. Some investigations based on local series or national registries aimed to evaluate the predictors and outcomes of major bleeding in TAVR patients^{7,10}. Peripheral vascular disease, end-stage renal disease, and coagulopathy have been identified to increase the rates of major bleeding.

Against this background, we aimed to stratify patients from the SCOPE 2 trial¹¹, a randomised comparison of the ACURATE *neo* (Boston Scientific) and the CoreValve Evolut (Medtronic) in 796 high-risk TAVR patients, according to their bleeding risk status based on the ARC-HBR definitions and to compare an array of key clinical outcomes including Bleeding Academic Research Consortium (BARC) 3-5 type bleeding complications in ARC-HBR positive (HBR+) and ARC-HBR negative patients (HBR-).

Editorial, see page 448

Methods

STUDY DESIGN

The SCOPE 2 trial design and rationale along with the principal 1-year results have been reported previously¹¹. In brief, SCOPE 2 was a multicentre, randomised, parallel-design, non-inferiority,

open-label trial carried out in 23 tertiary heart valve centres in 6 countries. It compared the safety and effectiveness of transfemoral TAVR using the ACURATE *neo* versus the CoreValve Evolut in patients with symptomatic severe aortic stenosis deemed to be at increased risk for mortality with SAVR, as assessed by the local Heart Teams. Inclusion criteria were the presence of symptomatic aortic stenosis with aortic annulus diameters covered by the sizes of the ACURATE *neo* and CoreValve Evolut valves. Left ventricular ejection fraction (<20%), pre-existing prosthetic valves in the aortic and/or mitral positions, bicuspid or unicuspid valves, severe mitral regurgitation, and/or peripheral anatomy inappropriate for transfemoral implant due to size, disease and degree of calcification or tortuosity of the aorta or ilio-femoral arteries represented the main criteria for exclusion. Complete details of the inclusion and exclusion criteria have been reported¹¹. The trial was conducted according to the Declaration of Helsinki and Good Clinical Practice and was approved by the investigational review board or research ethics committee at each participating centre. All participants gave their informed consent. Procedural recommendations were per standard of care. The mode of anaesthesia was selected according to local standard practice. Pre- and post-dilatation were performed at the operator's discretion, although predilatation is recommended by the manufacturer of the ACURATE *neo* valve. Access site closure was performed according to local practice. Minimally required laboratory analyses included haemoglobin, creatinine, and high-sensitivity troponin values. Dual antiplatelet therapy (preferably with aspirin and clopidogrel) was recommended for at least 3 months, followed by single antiplatelet therapy. In patients with an indication for oral anticoagulation or who had undergone recent coronary stent implantation, combination regimens and their duration were given at the discretion of the operator. Clinical follow-up was performed at 30 days and 1 year.

ARC-HBR CRITERIA

Some of the ARC-HBR criteria needed to be modified or were not available because they were either not captured in the electronic data capture or represented criteria for exclusion in the trial, as summarised in **Supplementary Table 1**. This approach has been followed in other validation studies¹²⁻¹⁵. Major and minor ARC-HBR criteria applied in the current study are as follows: age ≥ 75 years (minor); oral anticoagulant or novel oral anticoagulant at discharge (major); estimated glomerular filtration rate (eGFR) < 30 ml/min (major) and $eGFR \geq 30, < 60$ ml/min (minor); baseline haemoglobin < 11 g/dL (major), and 11-12.9 g/dL for men and 11-11.9 g/dL for women (minor); thrombocytes at index procedure $< 100 \times 10^9/L$ (major); non-steroidal anti-inflammatory drugs (NSAID) at discharge (minor); active cancer in the past 12 months (major); previous intracranial bleeding (major); any ischaemic stroke at any time not meeting the major criterion (minor). Patients were at HBR if at least one major criterion or two minor criteria were met¹⁰. An overall ARC-HBR score was calculated by adding 1 point for any major criterion and 0.5 for any minor criterion.

OUTCOMES

The primary endpoint was major or life-threatening bleeding (BARC type 3 or 5) at 12 months¹². Other key clinical outcomes were the occurrence of death, stroke, hospitalisation for valve-related symptoms or worsened chronic heart failure, myocardial infarction, new permanent pacemaker implantation, and any arrhythmia responsible for haemodynamic disorders at 30 days and at 1 year. All patients were followed up to 12 months. The definitions of all endpoints have been reported previously¹⁰. An independent committee adjudicated events after a review of original source documents.

STATISTICAL ANALYSIS

Discrete variables are expressed as percentages with frequencies and were compared by the χ^2 or Fisher's exact tests. Continuous variables are reported as mean \pm standard deviation and were compared by t-test if normally distributed or the Wilcoxon rank sum test for a non-parametric distribution. Event rates were based on Kaplan-Meier estimates in time-to-first-event analyses if mortality was part of the endpoint, or on cumulative incidence functions with the delta method for the estimation of the standard error, taking mortality as competing risk into account otherwise. Hazard ratios (HR) with 95% confidence intervals (CI) were determined by Cox regression analysis, and event rates were compared with the log-rank or Gray's test, respectively. The day of the procedure was taken as day 0. For patients without a procedure, the day of randomisation was taken as day 0. Interaction testing was performed to determine whether the relative risk of BARC 3 or 5 type bleeding and mortality measures at 12 months varied by age, number of anti-thrombotic medications, Society of Thoracic Surgeons (STS) score, and presence of oral anticoagulants at the time of the procedure. A landmark analysis was performed for BARC 3 or 5 type bleeding from 30 days to 12 months of follow-up according to presence or absence of oral anticoagulants (OAC) at day 30. Wolbers's adaptation of Harrell's C-statistic for survival data was used to describe the prediction accuracy of HBR for major bleeds¹⁶. The C-index was 0.52. All analyses were performed in the intention-to-treat population. A two-sided p-value <0.05 was considered significant. All statistical analyses were performed with SAS software (Version 9.4, SAS/STAT version 15.1; SAS Institute Inc.).

Results

Between April 2017 and April 2019, 796 patients with symptomatic severe aortic stenosis were randomised to ACURATE *neo* (n=398) versus CoreValve Evolut (n=398). Due to missing values, nine patients could not be classified as high or low bleeding risk patients. Of the remaining 787 patients, 633 patients (80.4%) were stratified as high bleeding risk (HBR+) according to the ARC-HBR criteria.

BASELINE CHARACTERISTICS

The main baseline characteristics of the study population are shown in **Table 1**. In brief, HBR+ patients were older and carried

Table 1. Baseline characteristics of the patients stratified by ARC-HBR status.

	HBR+ N=663	HBR- N=154	p-value
Gender, female	420 (66.3%)	113 (73.4%)	0.09
Age (yrs)	83.5 \pm 4.3	82.1 \pm 4.0	<0.0001
BMI (kg/m ²)	26.4 \pm 5.0	27.5 \pm 5.0	0.37
Symptoms			
NYHA Class III or IV	424 (67.0%)	87 (56.5%)	0.02
CCS Class III or IV	30 (4.7%)	10 (6.5%)	0.41
Syncope	72 (11.4%)	19 (12.3%)	0.78
STS predicted risk of mortality (%)	4.9 \pm 2.9	3.3 \pm 2.1	<0.0001
Medical conditions and medical history			
Diabetes	189 (29.9%)	32 (20.8%)	0.028
Dyslipidaemia	326 (51.5%)	78 (50.7%)	0.86
Hypertension	547 (86.4%)	129 (83.8%)	0.44
Current smoker	22 (3.5%)	5 (3.3%)	>0.99
Coronary artery disease	267 (42.2%)	52 (33.7%)	0.07
COPD	77 (12.2%)	15 (9.7%)	0.49
Extracranial cerebral artery disease	33 (5.2%)	7 (4.6%)	0.84
Peripheral artery disease	61 (9.6%)	10 (6.5%)	0.27
Dialysis	5 (0.8%)	0 (0.0%)	0.59
History of atrial fibrillation	252 (39.8%)	12 (7.8%)	<0.001
Previous pacemaker implantation	66 (10.4%)	8 (5.2%)	0.046
History of myocardial infarction	54 (8.5%)	12 (7.8%)	0.87
History of PCI	164 (25.9%)	38 (24.7%)	0.84
History of cardiac surgery	40 (6.3%)	4 (2.6%)	0.08
Previous aortic valvuloplasty	11 (1.7%)	1 (0.6%)	0.48
Prior cerebrovascular accident	99 (15.6%)	0 (0.0%)	<0.001
CT findings			
Aortic annulus perimeter (mm)	74 \pm 5	74 \pm 5	0.18
Aortic annulus area (mm ²)	423 \pm 56	420 \pm 55	0.50
Area-derived diameter (mm)	23 (22-24)	23 (22-25)	0.98
Number of events (percentages). Mean \pm standard deviation or median (Q1-Q3). BMI: body mass index; CCS: Canadian Cardiovascular Society; COPD: chronic obstructive pulmonary disease; CT: computed tomography; HBR: high bleeding risk; NYHA: New York Heart Association; PCI: percutaneous coronary intervention; STS: Society of Thoracic Surgeons			

less favourable medical conditions. Compared to HBR- patients, they were more frequently diabetic and more frequently had a history of permanent pacemaker implantation, atrial fibrillation and/or a prior cerebrovascular accident. Their estimated STS predicted risk of mortality (STS-PROM) at 30 days was higher (4.9% \pm 2.9% vs 3.3% \pm 2.1%; p<0.0001). The distribution of medical conditions leading to HBR+ status is depicted in **Figure 1**. HBR+ patients were more frequently on OAC therapy (vitamin K antagonists [VKA]) or novel oral anticoagulants (NOAC), but the total number of antithrombotic medications was similar between groups (**Table 2**).




Figure 1. Distribution of medical conditions leading to HBR+ status. CVA: cardiovascular accident; eGFR: estimated glomerular filtration rate; F/M: female/male; HBR: high bleeding risk; NOAC: novel oral anticoagulants; TIA: transient ischaemic attack

PROCEDURAL CHARACTERISTICS AND IMMEDIATE OUTCOME

The procedural characteristics did not vary significantly between HBR+ and HBR- patients (**Table 3**). A suture-mediated closure device was used in >80% of the patients. The immediate complication rates were similar between the two groups (13.4% vs 13.3%; $p>0.99$).

OUTCOMES AT 30 DAYS AND 12 MONTHS

Estimates of clinical outcomes at 30-day follow-up are shown in **Table 4**. BARC 3 or 5 type bleeding did not differ significantly between HBR+ and HBR- patients. HBR+ patients had higher rates of composite all-cause death and disabling stroke (4.8% vs 1.3%, risk difference 3.50%; 95% CI: 1.00-5.99; $p=0.0061$).

At 12 months, the rates of all-cause death were higher in HBR+ compared to HBR- patients (**Table 5**). However, the rates of BARC 3 or 5 type bleeding were not statistically different between groups (7.7% vs 6.1%, risk difference 1.67%; 95% CI: -2.72 to 6.06; $p=0.46$) (**Central illustration, Figure 2**). Compared to HBR- patients, HBR+ had more frequent access site-related BARC 3 or 5 type bleeding (3.0% vs. 0.7%, risk difference 2.36%; 95% CI: 0.49-4.23; $p=0.0133$). However, the rates of non-access site-related BARC 3 or 5 type bleeding were not statistically different between groups (4.6% vs 4.8% for HBR+ and HBR- patients, respectively, risk difference -0.17%; 95% CI: -3.99 to -3.65; $p=0.93$).

Subgroup analyses for bleeding events showed no significant interaction with patients' age (**Figure 3**). In addition, a landmark analysis conducted according to the presence or absence of OAC at day 30 to determine the risk of BARC 3-5 bleeding up to one year showed no difference between HBR+ and HBR- patients (**Table 6**). In the subgroup analysis for mortality at 12 months (**Figure 4**) a significant difference for intermediate-risk (STS 5-8)

Table 2. Anti-thrombotic medications at baseline, 30-day and 1-year follow-up, by HBR status.

		HBR+	HBR-
At baseline			
Aspirin		305/633 (48.2%)	99/154 (64.3%)
Clopidogrel		124/633 (19.6%)	29/154 (18.8%)
Prasugrel		2/633 (0.3%)	0/154 (0.0%)
Ticagrelor		6/633 (0.9%)	0/154 (0.0%)
Vitamin K antagonist		91/633 (14.4%)	0/154 (0.0%)
NOAC		142/633 (22.4%)	0/154 (0.0%)
No of antithrombotic medications	0	111/633 (17.5%)	52/154 (33.8%)
	1	389/633 (61.5%)	76/154 (49.4%)
	2	118/633 (18.6%)	26/154 (16.9%)
	3	15/633 (2.4%)	0/154 (0.0%)
At day 30			
Aspirin		334/573 (58.3%)	126/146 (86.3%)
Clopidogrel		252/573 (44.0%)	73/146 (50.0%)
Prasugrel		1/573 (0.2%)	0/146 (0.0%)
Ticagrelor		4/573 (0.7%)	0/146 (0.0%)
Vitamin K antagonist		69/573 (12.0%)	4/146 (2.7%)
NOAC		178/573 (31.1%)	10/146 (6.9%)
No of antithrombotic medications	0	15/573 (2.6%)	4/146 (2.7%)
	1	291/573 (50.8%)	71/146 (48.6%)
	2	254/573 (44.3%)	71/146 (48.6%)
	3	13/573 (2.3%)	0/146 (0.0%)
At 1 year			
Aspirin		255/504 (50.6%)	109/137 (79.6%)
Clopidogrel		78/504 (15.5%)	20/137 (14.6%)
Prasugrel		0/504 (0.0%)	0/137 (0.0%)
Ticagrelor		0/504 (0.0%)	0/137 (0.0%)
Vitamin K antagonist		55/504 (10.9%)	4/137 (2.9%)
NOAC		165/504 (32.7%)	17/137 (12.4%)
No of antithrombotic medications	0	26/504 (5.2%)	4/137 (2.9%)
	1	404/504 (80.2%)	116/137 (84.7%)
	2	73/504 (14.5%)	17/137 (12.4%)
	3	1/504 (0.2%)	0/137 (0.0%)
Number of events (percentages). NOAC: novel oral anticoagulants			

and older patients (81-85 and >85 years) was observed. Patients with BARC 3 or 5 type bleeding had a 27.4% rate of death at 12 months, an almost three-fold increase ($p<0.0052$) compared to patients who did not experience BARC 3 or 5 type bleeding during follow-up (9.6%). Further, the rate of death at 12 months in HBR+ patients was 30.9% after BARC type 3 or 5 bleeding and 11.0% in the absence of severe bleeding events. In HBR- patients, it was 11.1% and 3.9%, respectively.

Discussion

The present analysis of the SCOPE 2 trial indicates that HBR+ patients, identified based on ARC-HBR standards for PCI, had

Table 3. Procedure characteristics and outcomes by ARC-HBR status.


	HBR+ N=633	HBR- N=154	p-value
Procedure performed	620/633 (97.9%)	151/154 (98.1%)	>0.99
Total procedure time (min)	72±36	77±37	0.18
Total contrast volume administered (mL)	132±53	135±70	0.51
General anaesthesia	88/620 (14.2%)	28/151 (18.5%)	0.20
Type of access			0.39
Percutaneous	616/619 (99.5%)	151/151 (100.0%)	
Surgical cut-down	3/619 (0.5%)	0/151 (0.0%)	
Sheathless	235/620 (37.9%)	52/151 (34.4%)	0.45
Suture device closure for main access site			0.92
No	5/619 (0.8%)	1/151 (0.7%)	
ProStar	131/619 (21.2%)	29/151 (19.2%)	
1 ProGlide	34/619 (5.5%)	7/151 (4.6%)	
2 ProGlide	335/619 (54.1%)	88/151 (58.3%)	
Other	114/619 (18.4%)	26/151 (17.2%)	
Predilatation balloon valvuloplasty	370/620 (59.7%)	93/151 (61.6%)	0.71
Size device (mm)	27±2	26±2	0.08
Post-dilatation	248/620 (40.0%)	68/151 (45.0%)	0.27
Procedural complications	84/620 (13.5%)	28/151 (13.2%)	>0.99
Valve malpositioning	5/620 (0.8%)	6/151 (4.0%)	0.01
Coronary artery obstruction	2/620 (0.3%)	0/151 (0.0%)	>0.99
Haemodynamic instability	9/620 (1.5%)	0/151 (0.0%)	0.21
Cardiac tamponade	7/620 (1.1%)	1/151 (0.7%)	>0.99
Annular rupture	1/620 (0.2%)	1/151 (0.7%)	0.35
Conversion to open heart surgery	2/620 (0.3%)	0/151 (0.0%)	>0.99
Access vessel complication	48/620 (7.7%)	9/151 (6.0%)	0.60
Bleeding	16/620 (2.6%)	1/151 (0.7%)	0.22
Intraprocedural death	3/620 (0.5%)	0/151 (0.0%)	>0.99

Number of events (percentages). ARC-HBR: Academic Research Consortium - High Bleeding Risk; min: minutes; mL: millilitres; mm: millimetres

higher rates of death and stroke at 12 months, but similar rates of BARC 3 or 5 type bleeding compared to HBR- patients. These findings suggest that ARC-HBR criteria defined in a PCI

population are not relevant to discriminate an increased bleeding risk in TAVR patients. A contributing role of age is likely, as TAVR patients are on average 15 to 20 years older than patients

EuroIntervention

CENTRAL ILLUSTRATION Subgroup analysis of the SCOPE 2 trial population according to ARC-HBR criteria.

A-fib: atrial fibrillation; ARC: Academic Research Consortium; CI: confidence interval; HBR: high bleeding risk; OAC: oral anticoagulants; STS-PROM: Society of Thoracic Surgeons predicted risk of mortality




Figure 2. Cumulative incidence curves. A) BARC 3-5 bleeding and B) freedom from all-cause mortality in HBR+ and in HBR- patients. BARC: Bleeding Academic Research Consortium; CI: confidence interval; HBR: high bleeding risk

undergoing PCI. An age cut-off of 75 years, as defined by the ARC-HBR definitions for PCI to indicate a minor criterion, is probably not adequate for TAVR patients. Indeed, the rate of major or life-threatening bleeding was 3.6% in PARTNER 3 (mean age 73.3±5.8 years) and 10.4% in PARTNER II, where

patients were older (81.5±6.7 years)^{3,4}. Factors that increase the risk of bleeding after TAVR include a high prevalence of chronic kidney disease, liver disease, peripheral vasculopathy, acquired thrombocytopenia, colonic malignancy, and acquired reversible von Willebrand factor deficiency¹⁷⁻²¹. Since PCI is

Table 4. Endpoints at 30 days by ARC-HBR status.

	HBR+ N=633	HBR- N=154	Risk difference [95% CI]	p-value
All-cause death	19 (3.1%) [2.0-4.8]	0 (0.0%) [NC-NC]	3.08 [NC-NC]	NC
Cardiac death	14 (2.3%) [1.4-3.8]	0 (0.0%) [NC-NC]	2.29 [NC-NC]	NC
Non-cardiac death	5 (0.8%) [0.3-1.9]	0 (0.0%) [NC-NC]	0.81 [NC-NC]	NC
All strokes*	27 (4.3%) [2.9-6.1]	3 (2.0%) [0.5-5.3]	2.35 [-0.40-5.09]	0.09
Disabling strokes*	14 (2.2%) [1.3-3.6]	2 (1.3%) [0.3-4.3]	0.90 [-1.28-3.08]	0.42
Non-disabling strokes*	13 (2.1%) [1.2-3.5]	1 (0.6%) [0.1-3.3]	1.45 [-0.25-3.14]	0.09
All-cause death and disabling strokes	30 (4.8%) [3.4-6.8]	2 (1.3%) [0.3-5.2]	3.50 [1.00-5.99]	0.0061
All-cause death and all strokes	43 (6.9%) [5.2-9.2]	3 (2.0%) [0.6-6.0]	4.94 [1.95-7.94]	0.0012
Hospitalisation for valve-related symptoms or worsened CHF*	13 (2.1%) [1.2-3.5]	1 (0.7%) [0.1-3.4]	1.46 [-0.27-3.19]	0.09
Life-threatening major bleeding (BARC type 3 or 5)*	39 (6.2%) [4.5-8.3]	5 (3.3%) [1.2-7.0]	2.94 [-0.47-6.34]	0.09
Life-threatening major bleeding (BARC type 3b or 5)*	15 (2.4%) [1.4-3.8]	2 (1.3%) [0.3-4.3]	1.09 [-1.10-3.27]	0.33
Myocardial infarction*	2 (0.3%) [0.1-1.1]	0 (0.0%) [NC-NC]	0.32 [NC-NC]	NC
Valve-related dysfunction requiring repeat procedure*	2 (0.3%) [0.1-1.1]	0 (0.0%) [NC-NC]	0.32 [NC-NC]	NC
Implantation of multiple valves*	1 (0.2%) [0.0-0.9]	4 (2.6%) [0.9-6.1]	-2.44 [-4.97-0.09]	0.06
New LBBB*	92 (14.6%) [12.0-17.5]	30 (19.7%) [13.8-26.4]	-5.06 [-12.0-1.84]	0.15
New permanent pacemaker implantation*	95 (15.2%) [12.5-18.2]	16 (10.6%) [6.3-16.1]	4.64 [-1.02-10.31]	0.11
Any arrhythmia resulting in haemodynamic instability or requiring therapy	30 (4.8%) [3.3-6.7]	3 (2.0%) [0.5-5.3]	2.79 [-0.01-5.60]	0.051

Number of events (percentages) [95% CI: confidence interval]. Percentages are Kaplan-Meier estimates or cumulative incidence estimates (indicated by *) taking mortality as a competing risk into account. BARC: Bleeding Academic Research Consortium; CHF: congestive heart failure; HBR: high bleeding risk; LBBB: left bundle branch block; NC: not calculated

Table 5. Endpoints at 1 year by ARC-HBR status.

	HBR+ N=633	HBR- N=154	Risk difference [95% CI]	p-value
All-cause death	73 (12.4%) [10.0-15.4]	6 (4.3%) [2.0-9.4]	8.09 [3.76-12.41]	0.0002
Cardiac death	42 (7.2%) [5.4-9.6]	3 (2.1%) [0.7-6.3]	5.12 [1.97-8.26]	0.0015
Non-cardiac death	31 (5.6%) [4.0-7.9]	3 (2.3%) [0.7-7.0]	3.33 [0.12-6.53]	0.0422
All strokes*	35 (5.7%) [4.1-7.7]	7 (4.7%) [2.1-9.0]	0.97 [-2.92-4.86]	0.62
Disabling strokes*	17 (2.8%) [1.7-4.3]	4 (2.5%) [0.9-6.3]	0.04 [-2.88-2.97]	0.98
Non-disabling strokes*	18 (3.0%) [1.8-4.5]	3 (2.0%) [0.6-5.4]	0.93 [-1.70-3.57]	0.49
All-cause death and disabling strokes	84 (14.1%) [11.6-17.2]	9 (6.3%) [3.3-11.9]	7.80 [2.89-12.71]	0.0019
All-cause death and all strokes	100 (16.8%) [14.0-20.0]	11 (7.5%) [4.2-13.2]	9.25 [4.02-14.47]	0.0005
Hospitalisation for valve-related symptoms or worsened CHF*	37 (6.3%) [4.5-8.4]	4 (2.8%) [0.9-6.5]	3.51 [0.20-6.83]	0.0380
Life-threatening major bleeding (BARC type 3 or 5)*	48 (7.7%) [5.8-10.0]	9 (6.1%) [3.0-10.7]	1.67 [-2.72-6.06]	0.46
Life-threatening major bleeding (BARC type 3b or 5)*	21 (3.4%) [2.2-5.1]	3 (2.0%) [0.6-5.3]	1.43 [-1.25-4.11]	0.30
Myocardial infarction*	9 (1.5%) [0.7-2.8]	0 (0.0%) [NC-NC]	1.51 [NC-NC]	NC
Valve-related dysfunction requiring repeat procedure*	4 (0.7%) [0.2-1.6]	0 (0.0%) [NC-NC]	0.66 [NC-NC]	NC
Endocarditis*	4 (0.7%) [0.2-1.7]	0 (0.0%) [NC-NC]	0.69 [NC-NC]	NC
New LBBB*	96 (15.4%) [12.7-18.3]	30 (19.7%) [13.8-26.4]	-4.32 [-11.3-2.61]	0.22
New permanent pacemaker implantation*	98 (15.8%) [13.0-18.7]	16 (10.6%) [6.3-16.1]	5.16 [-0.52-10.84]	0.07
Any arrhythmia resulting in haemodynamic instability or requiring therapy	37 (6.1%) [4.4-8.2]	4 (2.7%) [0.9-6.3]	3.40 [0.18-6.62]	0.0387

Number of events (percentages) [95% CI: confidence interval]. Percentages are Kaplan-Meier estimates or cumulative incidence estimates (indicated by *) taking mortality as a competing risk into account. ARC-HBR: Academic Research Consortium - High Bleeding Risk; BARC: Bleeding Academic Research Consortium; CHF: congestive heart failure; LBBB: left bundle branch block; NC: not calculated



**Figure 3. BARC 3-5 bleeding at one year by subgroups. CI: confidence interval; HBR: high bleeding risk; OAC: oral anticoagulants; NC: not calculated; STS: Society of Thoracic Surgeons**

Table 6. Landmark analysis according to OAC status at day 30 to assess the one-year BARC 3-5 bleeding risk.

OAC at day 30 (interaction $p=0.1461$)	HBR+	HBR-	Risk difference [95% CI]	p -value
No	6/322 (1.9%) [0.8-3.9]	1/132 (0.8%) [0.1-3.9]	1.11 [-1.04-3.26]	0.31
Yes	4/242 (1.7%) [0.6-4.0]	2/14 (14.9%) [2.4-37.8]	-13.2 [-32.3-5.97]	0.18

Number of events (percentages) [95% CI: confidence interval]. Percentages are cumulative incidence estimates taking mortality as a competing risk into account; CI: confidence interval. HBR: high bleeding risk; OAC: oral anticoagulants

**Figure 4. Mortality at one year by subgroups.** CI: confidence interval; HBR: high bleeding risk; OAC: oral anticoagulants; NC: not calculated; STS: Society of Thoracic Surgeons

most frequently performed via the radial approach with smaller sheaths and catheters, the presence of peripheral vasculopathy is notably absent from ARC-HBR criteria while the size, presence of calcifications, and tortuosity of the iliac and femoral arteries are strongly associated with TAVR-related vascular complications and bleeding. More recently, Navarese et al have developed a 6-item algorithm comprising blood haemoglobin and serum iron concentrations, oral anticoagulation and dual antiplatelet therapy, common femoral artery diameter, and creatinine clearance that was able to identify patients at high risk of bleeding within 30 days after TAVR¹⁹. The role of anti-thrombotic medications is crucial in determining the rate of bleeding in these patients. However, we found that despite having more frequent use of OAC and being stratified according to bleeding-associated conditions, HBR+ patients had similar rates of severe bleeding

compared to HBR-, which highlights the fact that important conditions reflecting the specificity of a TAVR population are missing in the ARC-HBR definition.

In the present study, approximately 7% of patients who underwent TAVR had BARC 3 or 5 type bleeding at one year, which is consistent with recent reports^{5,7}. A previous study reported that life-threatening bleeding after TAVR, as defined by the VARC criteria, occurred in approximately 15 to 20% of TAVR procedures²². Our estimate of major bleeding complications in TAVR was 7.2%, which we believe to be a more contemporary estimate and could be further ameliorated using a single antithrombotic agent (e.g., a VKA or an NOAC in patients with atrial fibrillation, or aspirin in patients without) if PCI is not concurrently performed⁴.

Major bleeding or vascular complications have decreased as TAVR technology evolves into smaller device and sheath sizes^{22,23},

and bleeding complications are inconsistent in the early literature²⁴. However, major bleeding is still associated with a three-fold increase in one-year mortality following TAVR⁵⁻⁷. Patients referred to TAVR are elderly, frail and at risk for both bleeding and ischaemic complications^{7,17,25}. Careful evaluation of risk and benefit is warranted to identify the optimal antithrombotic regimen, as major late bleeding complications are associated with an increased risk of mortality⁵. Our data support and strengthen a previous report²⁶ suggesting that ARC-HBR criteria are not suitable for TAVR patients. Further studies and initiatives are warranted to determine the conditions qualifying HBR for the specific subset of patients requiring percutaneous valvular interventions.

Limitations

The present analysis has several limitations. The present report describes a *post hoc* subgroup analysis according to bleeding risk in the SCOPE 2 trial. Outcomes were assessed among ARC-HBR+ and ARC-HBR- patients. As per trial protocol, the current subgroup analysis should be considered exploratory and hypothesis-generating as the trial did not meet its primary objective and was not powered for either comparative assessment of bleedings nor for this patient stratification. In addition, some of the ARC-HBR criteria were not captured in the trial. Most uncaptured medical conditions (cirrhosis and all severe coagulation conditions, active cancer with bad prognosis) were criteria for exclusion in the SCOPE 2 trial, and some of them (brain malformations) are very rare in this TAVR population. In addition, ischaemic strokes and transient ischaemic attacks were classified as minor criteria in the absence of timing of the events. We assumed that this would not significantly impact the determination of the patients' groups.

Conclusions

ARC-HBR criteria defined for PCI patients did not identify a subset of TAVR patients at increased rates of BARC 3 or 5 bleeding in the SCOPE 2 trial. Specific HBR criteria should be defined for TAVR patients. These findings are clinically relevant and have potential important implications, especially for the selection of post-TAVR antithrombotic regimens based on individual bleeding risk profiles.

Impact on daily practice

Severe bleeding after TAVR is associated with increased morbidity and mortality. As opposed to PCI-related risks of bleeding that have been defined by an ARC initiative, the conditions leading to severe bleeding after TAVR remain insufficiently explored. The present report confirms and strengthens the fact that conditions stratifying high bleeding risk criteria in a PCI population are not relevant to qualify high bleeding risk in the subset of valvular patients requiring percutaneous intervention. High bleeding risk criteria should be defined in a way that is specific to TAVR patients. A dedicated initiative is warranted.

Appendix. Authors' affiliations

1. Institut Cardiovasculaire Paris-Sud, Hôpital Privé Jacques Cartier, Ramsay-Santé, Massy, France; 2. Division of Cardiology Azienda Ospedaliero Universitaria "Policlinico-Vittorio Emanuele" University of Catania, Catania, Italy; 3. Department of Thoracic and Cardiovascular Surgery, Heart and Diabetes Center Northrhine-Westfalia, University Hospital, Ruhr-University Bochum, Bad Oeynhausen, Germany; 4. Department of Cardiology, Leipzig Heart Center, University of Leipzig, Leipzig, Germany; 5. Department of Interventional Cardiology, Heart and Diabetes Center North Rhine Westfalia, Bad Oeynhausen, Germany; 6. Department of Cardiology, Brighton & Sussex University Hospitals NHS Trust, Brighton, United Kingdom; 7. Department of Cardiology, Leeds General Infirmary, Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom; 8. Department of Interventional Cardiology, Elisabeth Hospital Essen, Essen, Germany; 9. Department of Cardio-Thoracic-Vascular diseases and transplantation, Azienda Ospedaliero-Universitaria Policlinico "G. Rodolico-San Marco", Catania, Italy; 10. Groupe CardioVasculaire Interventionnel, Clinique Pasteur, Toulouse, France; 11. Department of Cardiovascular Medicine, Humanitas Clinical and Research Center, Milano, Italy; 12. Department of Cardiology, Brest University Hospital, Brest, France; 13. Cardiology Department, IRCCS Policlinico San Donato, Milano, Italy; 14. Department of Cardiology, Lille University Hospital, Lille, France; 15. Department of Cardiology, Goethe University Hospital Frankfurt, Frankfurt am Main, Germany; 16. Interventional Cardiology Unit, IRCCS San Raffaele Scientific Institute, Milan, Italy; 17. KU Leuven, Faculty of Medicine, I-BioStat, Leuven, Belgium and UHasselt, I-BioStat, Hasselt, Belgium; 18. Department of Internal Medicine II, Medical University of Vienna, Vienna, Austria

Acknowledgments

The authors are grateful to Mrs. Dupic for her skillful assistance in reviewing the present manuscript.

Guest editor

This paper was guest edited by Franz-Josef Neumann, MD; Department of Cardiology and Angiology II, University Heart Center Freiburg - Bad Krozingen, Bad Krozingen, Germany.

Funding

The SCOPE 2 Trial is an investigator-initiated trial, funded by CERIC.

Conflict of interest statement

P. Garot is the medical Director and a shareholder of CERC. M-C. Morice is the CEO and a shareholder of CERC and a minor shareholder of Electroducer. C. Tamburino has received speaker fees from Medtronic. S. Bleiziffer has received speaker fees from Boston Scientific and Medtronic. M. Cunnington has received speaker and consultant fees from Medtronic. A. Wolf is a proctor for Boston Scientific, Edwards Lifesciences, and Medtronic. M. Barbanti has received consultant fees from Edwards Lifesciences. P. Pagnotta

is a proctor for Boston Scientific, Cardia, and Gore Medical. F. Bedogni has received personal fees from Abbott Vascular, Boston Scientific, Medtronic, Meril Life Sciences, and Terumo. E. Van Belle has received personal fees from Philips/Volcano. M. Vasa-Nicotera is a proctor for Boston Scientific and Medtronic. A. Chieffo has received speaker fees from Abiomed, Abbott Vascular, Biosensors, Cardinal Health, and Magenta, and consultant fees from Abiomed, Abbott Vascular, Biosensors, Cardinal Health, and Magenta. K. Bogaerts has received consultant fees from the Cardiovascular European Research Center. C. Hengstenberg is a proctor for Boston Scientific; reports institutional grants from Boston Scientific; has received speaker fees from Boston Scientific; and has received consultant fees from Boston Scientific. D. Capodanno has received speaker fees from AstraZeneca, Biosensors, Boehringer Ingelheim, Boston Scientific, Daiichi Sankyo, Menarini, Pfizer, and Sanofi, and consultant fees from Abbott Vascular, Amgen, AstraZeneca, and Bayer. The other authors have no conflicts of interest to declare. The Guest Editor reports lecture fees paid to his institution from Amgen, Bayer Healthcare, Biotronik, Boehringer Ingelheim, Boston Scientific, Daiichi Sankyo, Edwards Lifesciences, Ferrer, Pfizer, and Novartis; consultancy fees paid to his institution from Boehringer Ingelheim; and grant support from Bayer Healthcare, Boston Scientific, Biotronik, Edwards Lifesciences, GlaxoSmithKline, Medtronic, and Pfizer.

References

- Mack MJ, Leon MB, Thourani VH, Makkar R, Kodali SK, Russo M, Kapadia SR, Malaisrie SC, Cohen DJ, Pibarot P, Leipsic J, Hahn RT, Blanke P, Williams MR, McCabe JM, Brown DL, Babaliaros V, Goldman S, Szeto WY, Genereux P, Pershad A, Pocock SJ, Alu MC, Webb JG, Smith CR; PARTNER 3 Investigators. Transcatheter Aortic-Valve Replacement with a Balloon Expandable Valve in Low-Risk Patients. *N Engl J Med*. 2019;380:1695-705.
- Leon MB, Smith CR, Mack MJ, Makkar RR, Svensson LG, Kodali SK, Thourani VH, Tuzzu EM, Miller DC, Herrmann HC, Doshi D, Cohen DJ, Pichard AD, Kapadia S, Dewey T, Babaliaros V, Szeto WY, Williams MR, Kereiakes D, Zajarias A, Greason KL, Whisenant BK, Hodson RW, Moses JW, Trento A, Brown DL, Fearon WF, Pibarot P, Hahn RT, Jaber WA, Anderson WN, Alu MC, Webb JG; PARTNER 2 Investigators. Transcatheter or Surgical Aortic Valve Replacement in Intermediate-Risk Patients. *N Engl J Med*. 2016;374:1609-20.
- Otto CM, Nishimura RA, Bonow RO, Carabello BA, Erwin JP 3rd, Gentile F, Jneid H, Krieger EV, Mack M, McLeod C, O'Gara PT, Rigolin VH, Sundt TM 3rd, Thompson A, Toly C. 2020 ACC/AHA Guideline for the Management of Patients With Valvular Heart Disease: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2021;143:e35-71.
- Vahanian A, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J, Capodanno D, Conradi L, De Bonis M, De Paulis R, Delgado V, Freemantle N, Haugaa KH, Jeppsson A, Jüni P, Pierard L, Prendergast BD, Sadaba JR, Tribouilloy C, Wojakowski W. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *EuroIntervention*. 2022;17:e1126-96.
- Bendayan M, Messas N, Perrault LP, Asgar AW, Lauck S, Kim DH, Arora RC, Langlois Y, Piazza N, Martucci G, Lefèvre T, Noiseux N, Lamy A, Peterson MD, Labinaz M, Popma JJ, Webb JG, Afilalo J. Frailty and Bleeding in Older Adults Undergoing TAVR or SAVR: Insights From the FRAILTY-AVR Study. *JACC Cardiovasc Interv*. 2020;13:1058-68.
- Ullah W, Jafar M, Zahid S, Ahmed F, Khan MZ, Sattar Y, Fischman DL, Virani SS, Alam M. Predictors of In-Hospital Mortality in Patients With End-Stage Renal Disease Undergoing Transcatheter Aortic Valve Replacement: A Nationwide Inpatient Sample Database Analysis. *Cardiovasc Revasc Med*. 2022;34:63-8.
- Khan H, Gilani A, Qayum I, Khattak T, Haq F, Zahid Anwar M, Khan MA, Asjad SJ, Abbas S, Inayat A. An Analysis of the Predictors of Major Bleeding After Transcatheter Aortic Valve Transplantation Using the National Inpatient Sample (2015-2018). *Cureus*. 2021;13:e16022.
- Urban P, Mehran R, Collieran R, Angiolillo DJ, Byrne RA, Capodanno D, Cuisset T, Cutlip D, Eerdmans P, Eikelboom J, Farb A, Gibson CM, Gregson J, Haude M, James SK, Kim HS, Kimura T, Konishi A, Laschinger J, Leon MB, Magee PFA, Mitsutake Y, Mylotte D, Pocock S, Price MJ, Rao SV, Spitzer E, Stockbridge N, Valgimigli M, Varenne O, Windhoevel U, Yeh RW, Krucoff MW, Morice MC. Defining High Bleeding Risk in Patients Undergoing Percutaneous Coronary Intervention. *Circulation*. 2019;140:240-61.
- VARC-3 WRITING COMMITTEE; Généreux P, Piazza N, Alu MC, Nazif T, Hahn RT, Pibarot P, Bax JJ, Leipsic JA, Blanke P, Blackstone EH, Finn MT, Kapadia S, Linke A, Mack MJ, Makkar R, Mehran R, Popma JJ, Reardon M, Rodes-Cabau J, Van Mieghem NM, Webb JG, Cohen DJ, Leon MB. Valve Academic Research Consortium 3: Updated Endpoint Definitions for Aortic Valve Clinical Research. *J Am Coll Cardiol*. 2021;77:2717-46.
- Tchetché D, Van der Boon RM, Dumonteil N, Chieffo A, Van Mieghem NM, Farah B, Buchanan GL, Saady R, Marcheix B, Serruys PW, Colombo A, Carrie D, De Jaegere PP, Fajadet J. Adverse impact of bleeding and transfusion on the outcome post-transcatheter aortic valve implantation: insights from the Pooled-Rotterdam-Milano-Toulouse In Collaboration Plus (PRAGMATIC Plus) initiative. *Am Heart J*. 2012;164:402-9.
- Tamburino C, Bleiziffer S, Thiele H, Scholtz S, Hildick-Smith D, Cunningham M, Wolf A, Barbanti M, Tchetché D, Garot P, Pagnotta P, Gilard M, Bedogni F, Van Belle E, Vasa-Nicotera M, Chieffo A, Deutsch O, Kempfert J, Sondergaard L, Butter C, Trillo-Nouche R, Lotfi S, Möllmann H, Joner M, Abdel-Wahab M, Bogaerts K, Hengstenberg C, Capodanno D. Comparison of Self-Expanding Bioprostheses for Transcatheter Aortic Valve Replacement in Patients With Symptomatic Severe Aortic Stenosis: SCOPE 2 Randomized Clinical Trial. *Circulation*. 2020;142:2431-42.
- Cao D, Mehran R, Dangas G, Baber U, Sartori S, Chandiraman R, Stefanini GG, Angiolillo DJ, Capodanno D, Urban P, Morice MC, Krucoff M, Goel R, Roumeliotis A, Sweeny J, Sharma SK, Kini A. Validation of the Academic Research Consortium High Bleeding Risk Definition in Contemporary PCI Patients. *J Am Coll Cardiol*. 2020;75:2711-22.
- Corpataux N, Spirito A, Gragnano F, Vaisnora L, Galea R, Svab S, Gargiulo G, Zanchin T, Zanchin C, Siontis GCM, Praz F, Lanz J, Hunziker L, Stortecky S, Pilgrim T, Räber L, Capodanno D, Urban P, Pocock S, Heg D, Windecker S, Valgimigli M. Validation of high bleeding risk criteria and definition as proposed by the academic research consortium for high bleeding risk. *Eur Heart J*. 2020;41:3743-9.
- Gragnano F, Spirito A, Corpataux N, Vaisnora L, Galea R, Gargiulo G, Siontis GCM, Praz F, Lanz J, Billinger M, Hunziker L, Stortecky S, Pilgrim T, Bär S, Ueki Y, Capodanno D, Urban P, Pocock SJ, Mehran R, Heg D, Windecker S, Räber L, Valgimigli M. Impact of Clinical Presentation on Bleeding Risk after Percutaneous Coronary Intervention and Implications for the ARC-HBR Definition. *EuroIntervention*. 2021;17:e898-909.
- Mehran R, Rao SV, Bhatt DL, Gibson CM, Caixeta A, Eikelboom J, Kaul S, Wiviott SD, Menon V, Nikolsky E, Serebruany V, Valgimigli M, Vranckx P, Taggart D, Sabik JF, Cutlip DE, Krucoff MW, Ohman EM, Steg PG, White H. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation*. 2011;123:2736-47.
- Wolbers M, Koller MT, Witteman JC, Steyerberg EW. Prognostic models with competing risks: methods and application to coronary risk prediction. *Epidemiology*. 2009;20:555-61.
- Ferlini M, Mauri S, Rossini R. Dual antiplatelet therapy after TAVR: a drop in the bucket? *Int J Cardiol*. 2019;280:46-8.
- Ullah W, Zahid S, Hamzeh I, Birnbaum Y, Virani SS, Alam M. Trends and Predictors of Transcatheter Aortic Valve Implantation Related In-Hospital Mortality (From the National Inpatient Sample Database). *Am J Cardiol*. 2021;143:97-103.
- Navarese EP, Zhang Z, Kubica J, Andreotti F, Farinaccio A, Bartorelli AL, Bedogni F, Ruppj M, Tomai F, Giordano A, Reimers B, Spaccarotella C, Wilczek K, Stepinska J, Witkowski A, Grygier M, Kukulski T, Wanha W, Wojakowski W, Lesiak M, Dudek D, Zembala MO, Berti S; a Joint Effort of the Italian and Polish Cardiac Interventional Societies. Development and Validation of a Practical Model to Identify Patients at Risk of Bleeding After TAVR. *JACC Cardiovasc Interv*. 2021;14:1196-206.
- Zahid S, Ullah W, Khan MU, Abbas S, Ud Din MT, Uddin MF, Inayat A, Ubaid A, Salman F, Thakkar S, Salama A, Khan MZ. Trends, predictors, and outcomes of major bleeding after transcatheter aortic valve implantation, from national inpatient sample (2011-2018). *Expert Rev Cardiovasc Ther*. 2021;19:557-63.
- Zhang X, Wang T, Lan R, Dai Q, Kang L, Wang L, Wang Y, Xu W, Xu B. Meta-Analysis Comparing Results of Transcatheter Versus Surgical Aortic-Valve Replacement in Patients With Severe Aortic Stenosis. *Am J Cardiol*. 2020;125:449-58.
- Kolte D, Vlahakes GJ, Palacios IF, Sakhuja R, Passeri JJ, Ingleessis I, Elmariah S. Transcatheter Versus Surgical Aortic Valve Replacement in Low-Risk Patients. *J Am Coll Cardiol*. 2019;74:1532-40.

23. Dangas GD, Tijssen JGP, Wohrle J, Sondergaard L, Gilard M, Mollmann H, Makkar RR, Herrmann HC, Giustino G, Baldus S, De Backer O, Guimaraes AHC, Gullestad L, Kini A, von Lewinski D, Mack M, Moreno R, Schafer U, Seeger J, Tchetché D, Thomitzek K, Valgimigli M, Vranckx P, Welsh RC, Wildgoose P, Volkl AA, Zazula A, van Amsterdam RGM, Mehran R, Windecker S, GALILEO Investigators. A controlled trial of rivaroxaban after transcatheter aortic-valve replacement. *N Engl J Med*. 2020;382:120-9.

24. Généreux P, Head SJ, Van Mieghem NM, Kodali S, Kirtane AJ, Xu K, Smith C, Serruys PW, Kappetein AP, Leon MB. Clinical outcomes after transcatheter aortic valve replacement using valve academic research consortium definitions: a weighted meta-analysis of 3,519 patients from 16 studies. *J Am Coll Cardiol*. 2012;59:2317-26.

25. Mangieri A, Montalto C, Poletti E, Sticchi A, Crimi G, Giannini F, Latib A, Capodanno D, Colombo A. Thrombotic Versus Bleeding Risk After Transcatheter Aortic Valve Replacement: JACC Review Topic of the Week. *J Am Coll Cardiol*. 2019;74:2088-101.

26. Bor WL, Chan Pin Yin DRPP, Brouwer J, Nijenhuis VJ, Peper J, Timmers L, Rensing BJWM, Swaans MJ, Ten Berg JM. Assessment of the Academic Research Consortium for High Bleeding Risk Criteria in Patients Undergoing TAVR. *JACC Cardiovasc Interv*. 2021;14:1265-7.

Supplementary data

Supplementary Table 1. Comparison of definitions between the ARC-HBR criteria and the present study.

The supplementary data are published online at:
[https://eurointervention.pcronline.com/
doi/10.4244/EIJ-D-21-01048](https://eurointervention.pcronline.com/doi/10.4244/EIJ-D-21-01048)



Supplementary data

Supplementary Table 1. Comparison of definitions between the ARC-HBR criteria and the present study.

ARC-HBR criteria	Present study	Category	Comments
Age ≥ 75	Age ≥ 75	Minor	Identical
Anticipated use of long-term oral anticoagulation*	Long-term OAC	Major	Identical
Severe or end-stage CKD (eGFR < 30 mL/min)	Clearance creatinine by original MDRD, same cut-off	Major	Identical
Moderate CKD (eGFR 30–59 mL/min)	Clearance creatinine by original MDRD, same cut-off	Minor	Identical
Haemoglobin < 11 g/dL	Haemoglobin < 11 g/dL	Major	Identical
Haemoglobin 11–12.9 g/dL for men and 11–11.9 g/dL for women	Haemoglobin 11–12.9 g/dL for men and 11–11.9 g/dL for women	Minor	Identical
Spontaneous bleeding requiring hospitalisation or transfusion in the past 6 months or at any time, if recurrent		Major	Not available
Spontaneous non-intracranial bleeding requiring hospitalisation or transfusion within the past 12 months not meeting the major criterion		Minor	Not available
Moderate or severe baseline thrombocytopaenia (platelet count $< 100 \times 10^9/L$)	Severe coagulation condition is a criterion for exclusion	Major	Not available
Chronic bleeding diathesis	Active bleeding is a criterion for exclusion	Major	Not available
Liver cirrhosis with portal hypertension	Severe coagulation condition is a criterion for exclusion	Major	Not available
Long-term use of oral NSAIDs or steroids NSAIDS at discharge	List of medications at the time of procedure and at discharge available	Minor	identical
Active malignancy (excluding non-melanoma skin cancer) within the past 12 months	Active cancer in the past 12 months	Major	identical
Previous spontaneous ICH (at any time)	Previous haemorrhagic stroke	Major	modified
Presence of a bAVM		Major	Not available

Moderate or severe ischaemic stroke within the past 6 mo	Previous stroke or TIA (no indication of date)	Major	Not available
Non-deferrable major surgery on DAPT		Major	Not available
Recent major surgery or major trauma within 30 d before PCI		Major	Not available
Any ischaemic stroke at any time not meeting the major criterion	Previous stroke or TIA (no indication of date)	Minor	Modified

bAVM: brain arteriovenous malformation; CKD: chronic kidney disease; DAPT: dual antiplatelet therapy; eGFR: estimated glomerular filtration rate; HBR: high bleeding risk; ICH: intracranial haemorrhage; NSAID: non-steroidal anti-inflammatory drug; PCI: percutaneous coronary intervention